

On existence of singular solutions

Miroslav Bartušek

Department of Mathematics, Masaryk University, Janáčkovo nám. 2a, Brno 662 95, Czech Republic

Received 29 April 2002

Submitted by J.S.W. Wong

Abstract

In the paper sufficient conditions are given under which the differential equation $y^{(n)} = f(t, y, \dots, y^{(n-2)})g(y^{(n-1)})$ has a singular solution $y: [T, \tau) \rightarrow R$, $\tau < \infty$ fulfilling

$$\lim_{t \rightarrow \tau} |y^{(j)}(t)| = \infty, \quad j = 0, 1, \dots, n-1.$$

© 2003 Elsevier Science (USA). All rights reserved.

Consider the n -order differential equation

$$y^{(n)} = f(t, y, y', \dots, y^{(n-2)})g(y^{(n-1)}), \quad (1)$$

where $n \geq 2$, $f \in C^0(R_+ \times R^{n-1})$, $g \in C^0(R)$, $R_+ = [0, \infty)$, $R = (-\infty, \infty)$.

A solution y defined on $[T, \tau) \subset R_+$ is called singular if $\tau < \infty$ and y cannot be defined for $t = \tau$. Note that in this case $\limsup_{t \rightarrow \tau} |y^{(n-1)}| = \infty$. A singular solution y is called nonoscillatory if there exists a left neighbourhood of τ in which $y \neq 0$.

The problem of the nonexistence of singular solutions of (1) is solved by Wintner's theorem (see [8] or [11]); the main assumption is

$$|f(t, x_1, \dots, x_{n-1})g(x_n)| \leq r(t)\omega\left(\sum_{i=1}^n |x_i|\right) \quad \text{on } R_+ \times R^n,$$

where $\omega \in C^0(R_+)$ is nondecreasing, $\omega(x) > 0$ for $x > 0$, $\int_1^\infty (1/\omega(x)) dx = \infty$ and $r \in C^0(R_+)$.

For the second order Emden–Fowler equation

$$y'' = r(t)|y|^\lambda \operatorname{sgn} y, \quad \lambda > 0, \quad (2)$$

where $r \leq 0$ on R_+ the nonexistence result is extended in [6,10].

E-mail address: bartusek@math.muni.cz.

As concern the existence of singular solutions of (1), fulfilling Cauchy initial conditions, the first results for (2) are obtained in [9] and they are generalized for (1) and its special cases, e.g., in [3,5,11]. Note that if $r(t) \leq 0$ on R_+ , there exists no theorem for the existence of singular solutions of (2), but an example of a function $r < 0$ is given in [4] for which (2) has a singular solution.

In this paper another problem will be studied—we will seek for singular solutions with prescribed asymptotics in the right-hand side point of the definition interval.

More precisely, the existence problem of a singular solution y of (1) fulfilling

$$\begin{aligned} \tau &\in (0, \infty), \quad l \in \{0, 1, \dots, n-2\}, \quad c_i \in R, \quad i = 0, 1, \dots, l, \\ \lim_{t \rightarrow \tau_-} y^{(i)}(t) &= c_i \quad \text{for } i = 0, 1, \dots, l, \\ \lim_{t \rightarrow \tau_-} |y^{(j)}(t)| &= \infty \quad \text{for } j = l+1, \dots, n-1, \end{aligned} \quad (3)$$

has been studied where y is defined in a left neighbourhood of τ . Sufficient (necessary) conditions for the existence of a solution y fulfilling (3) are given in [7] (case $n = 2$) and in [1,2] (case $n \geq 2$). Recall, that singular solutions fulfilling (3) are sometimes called black-hole solutions (see [7]).

In the present paper we extend our investigation to the case that is not obtained in (3). We will seek a singular solution y of (1) that fulfills

$$\tau \in (0, \infty), \quad \lim_{t \rightarrow \tau_-} |y^{(i)}(t)| = \infty \quad \text{for } i = 0, 1, \dots, n-1. \quad (4)$$

Note that if y is a singular solution fulfilling (4) then

$$\operatorname{sgn} y^{(k)}(t) = \operatorname{sgn} y^{(s)}(t) \quad \text{for } k, s \in \{1, \dots, n-1\},$$

holds in a left neighbourhood of τ .

The following theorem gives us sufficient conditions for the nonexistence of solutions fulfilling (4).

Theorem 1. Let $\tau \in (0, \infty)$ and I be a left neighbourhood of τ , $I \subset R_+$. Let $M \in (0, \infty)$ and $M_1 \in (0, \infty)$ be such that one of the following assumptions holds:

$$(i) \quad \alpha f(t, x_1, \dots, x_{n-1})g(x_n) \leq 0 \quad \text{for } \alpha x_i \geq M, \\ i = 1, 2, \dots, n, \quad \alpha \in \{-1, 1\}, \quad t \in I;$$

$$(ii) \quad \lambda > 1 + \frac{1}{n-1}$$

and

$$\alpha f(t, x_1, \dots, x_{n-1})g(x_n) \geq M_1 |x_n|^\lambda \quad \text{for } \alpha x_i \geq M, \\ i = 1, \dots, n, \quad \alpha \in \{-1, 1\}, \quad t \in I.$$

Then there exists no solution y of (1) fulfilling (4).

Proof. Let y be a solution of (1) defined in a left neighbourhood $J \subset I$ of τ such that $\lim_{t \rightarrow \tau_-} y^{(j)}(t) = \infty$, $j = 0, 1, \dots, n-1$ (the case $\lim_{t \rightarrow \tau_-} y^{(j)}(t) = -\infty$, $j = 0, 1, \dots, n-1$ can be investigated similarly). Put $\alpha = 1$. It is clear that there exists $\tau_1 \in J$ such that

$$y^{(j)}(t) \geq M \quad \text{for } t \in J_1 = [\tau_1, \tau), \quad i = 0, 1, \dots, n-1. \quad (5)$$

Let (i) be valid. Then (1) and (5) yield $y^{(n)}(t) \leq 0$ on J_1 that contradicts $\lim_{t \rightarrow \tau_-} y^{(n-1)} = \infty$ and $\tau < \infty$.

Let (ii) be valid. Put $\lambda_1 = 1/(\lambda - 1)$, hence $n - 1 - \lambda_1 > 0$. According to (1), (5) and the assumption (ii)

$$y^{(n)}(t) \geq M_1 [y^{(n-1)}(t)]^\lambda, \quad t \in J_1,$$

and the integration on $[t, \tau) \subset J_1$ yields

$$y^{(n-1)}(t) \leq [(\lambda - 1)M_1(\tau - t)]^{-\lambda_1}, \quad t \in J_1.$$

Hence, Taylor series theorem yields

$$\begin{aligned} \infty = y(\tau) &= \sum_{i=0}^{n-1} \frac{y^{(i)}(\tau_1)}{i!} (\tau - \tau_1)^i + \int_{\tau_1}^{\tau} \frac{(\tau - s)^{n-2}}{(n-2)!} y^{(n-1)}(s) ds \\ &\leq M_4 + M_5 \int_{\tau_1}^{\tau} (\tau - s)^{n-2-\lambda_1} ds < \infty, \end{aligned}$$

as $n - 2 - \lambda_1 > -1$; here

$$M_4 = \sum_{i=0}^{n-1} \frac{y^{(i)}(\tau_1)}{i!} (\tau - \tau_1)^i, \quad M_5 = \frac{[(\lambda - 1)M_1]^{-\lambda_1}}{(n-2)!}.$$

The contradiction proves that y does not fulfill (4). \square

To prove the existence of a singular solution fulfilling (4) we need the following lemma.

Lemma 1. Let $[a, b] \subset R_+$, $\Phi \in C^0[a, b]$ and $\tilde{f} \in C^0([a, b] \times R^n)$ be such that

$$\tilde{f}(t, x_1, \dots, x_n) \leq \Phi(t), \quad t \in [a, b], \quad x_i \in R, \quad i = 1, 2, \dots, n.$$

Then for arbitrary $c_i \in R$, $i = 0, 1, \dots, n-1$ the equation

$$u^{(n)} = \tilde{f}(t, u, \dots, u^{(n-1)})$$

has at least one solution fulfilling the boundary value conditions

$$u^{(i)}(a) = c_i, \quad i = 0, 1, \dots, n-2, \quad u(b) = c_{n-1}.$$

Proof. It follows, e.g., from [11, Lemma 10.1], as the homogeneous problem $u^{(n)} = 0$, $u^{(i)}(a) = 0$, $i = 0, 1, \dots, n-2$, $u(b) = 0$ has the trivial solution only. \square

The following theorem gives us a sufficient condition under which singular solutions with (4) exist.

Theorem 2. Let $\tau \in (0, \infty)$, $I = [\bar{\tau}, \tau] \subset R_+$, $\alpha \in \{-1, 1\}$, $M \in (0, \infty)$ and $M_1 \in (0, \infty)$ be such that

$$g(x) \geq |x|^\lambda \quad \text{for } \alpha x \geq M \quad (6)$$

and

$$\alpha f(t, x_1, \dots, x_{n-1}) \geq M_1 \quad \text{for } \alpha x_i \geq M, \quad i = 1, 2, \dots, n-1, \quad t \in I. \quad (7)$$

Let

$$1 < \lambda \leq 1 + \frac{1}{n-1}. \quad (8)$$

Then there exists a solution y of (1) defined in a left neighbourhood of $t = \tau$ that fulfills (4).

Proof. We prove the statement for $\alpha = 1$. For $\alpha = -1$ the proof is similar. Let $\lambda < 1 + 1/(n-1)$ and let

$$\begin{aligned} M_2 &= \max\{f(t, x_1, \dots, x_{n-1}) : t \in I, M \leq x_i \leq 3Me^\tau, i = 1, 2, \dots, n-1\}, \\ \lambda_1 &= \frac{1}{\lambda-1}, \quad M_3 = [M_1(\lambda-1)]^{-\lambda_1} + 2M\tau^{\lambda_1}, \quad k_0 > (n+1)M. \end{aligned} \quad (9)$$

Hence, according to (8)

$$n-1-\lambda_1 < 0. \quad (10)$$

Let $T \in I$, $T < \tau$ be such that

$$\tau - T \leq 1, \quad \int_M^{2M} \frac{ds}{g(s)} > M_2(\tau - T), \quad (11)$$

and denote $J = [T, \tau)$.

Consider an auxiliary two-point boundary-value problem: $k \in \{k_0, k_0 + 1, \dots\}$,

$$\begin{aligned} y^{(n)} &= f(t, \Phi_0(t, y), \dots, \Phi_0(t, y^{(n-2)}))g(\Phi(t, y^{(n-1)})), \\ y^{(i)}(T) &= M, \quad i = 0, 1, \dots, n-2, \quad \lim_{t \rightarrow \tau_-} y(t) = k, \end{aligned} \quad (12)$$

where

$$\begin{aligned} \varphi(t) &= M(n-1) + M_3[\lambda_1 - n + 1]^{-1}(\tau - t)^{-\lambda_1} \quad \text{if } n-1-\lambda_1 < 0, \\ \varphi(t) &= M(n-1) + M_3(\tau - t)^{-1} \quad \text{if } n-1-\lambda_1 = 0, \\ \Phi(t, s) &= \begin{cases} s & \text{for } M \leq s \leq M_3(\tau - t)^{-\lambda_1}, \\ M_3(\tau - t)^{-\lambda_1} & \text{for } s > M_3(\tau - t)^{-\lambda_1}, \\ M & \text{for } s < M, \end{cases} \\ \Phi_0(t, s) &= \begin{cases} s & \text{for } M \leq s \leq \varphi(t), \\ \varphi(t) & \text{for } s > \varphi(t), \\ M & \text{for } s < M. \end{cases} \end{aligned}$$

Note, that according to the definition of M_3 , Φ is well-defined.

To prove the existence of a solution of (12), let us consider the sequence of boundary value problems

$$\begin{aligned}\bar{m} &> \frac{1}{\tau - T}, \quad m \in \{\bar{m}, \bar{m} + 1, \dots\}, \quad \tau_m = \tau - \frac{1}{m}, \quad J_m = [T, \tau_m], \\ Z^{(n)} &= F(t, Z, Z', \dots, Z^{(n-1)}), \\ Z^{(i)}(T) &= M, \quad i = 0, 1, \dots, n-2, \quad Z(\tau_m) = k,\end{aligned}\tag{13}$$

where

$$F(t, x_1, \dots, x_n) = f(t, \Phi_0(t, x_1), \dots, \Phi_0(t, x_{n-1}))g(\Phi(t, x_n)).\tag{14}$$

As evidently $|F(t)|$ is bounded from above by a continuous function on J_m , Lemma 1 yields the existence of a solution Z_m of (13). It is clear that according to (6), (7) and (14)

$$Z_m^{(n)} > 0, \quad Z_m^{(n-1)} \text{ is increasing on } J_m.\tag{15}$$

Further, we estimate the derivatives of Z_m . First we prove that

$$M < Z_m^{(n-1)}(t), \quad t \in J_m, \quad m = \bar{m}, \bar{m} + 1, \dots\tag{16}$$

For this, we prove that

$$Z_m^{(n-1)}(\tau_m) > 2M.\tag{17}$$

Let, contrarily, $Z_m^{(n-1)}(\tau_m) \leq 2M$. Then the Taylor series theorem, (11) and (15) yield

$$\begin{aligned}k_0 \leq k = Z_m(\tau_m) &= M \sum_{i=0}^{n-2} \frac{(\tau_m - T)^i}{i!} + \int_T^{\tau_m} \frac{(\tau_m - s)^{n-2}}{(n-2)!} Z_m^{(n-1)}(s) ds \\ &\leq M(n-1) + 2M = (n+1)M,\end{aligned}$$

that contradicts (9). Hence (17) holds and if (16) is not valid there exist T_1 and T_2 such that $T \leq T_1 < T_2 < \tau_m$, $Z_m^{(n-1)}(T_1) = M$ and $Z_m^{(n-1)}(T_2) = 2M$. Then, according to (11), (15) and Taylor series theorem, we have

$$\begin{aligned}M \leq Z_m^{(i)}(t) &= M \sum_{j=0}^{n-i-2} \frac{(t - T_1)^j}{j!} + \int_{T_1}^t \frac{(t - s)^{n-i-2}}{(n-i-2)!} Z_m^{(n-1)}(s) ds \\ &\leq Me^\tau + 2M \int_{T_1}^t e^{t-s} ds \leq 3Me^\tau, \quad t \in J_m, \quad i = 0, 1, \dots, n-2.\end{aligned}\tag{18}$$

It follows from the definition of M_3 that $2M \leq M_3(\tau - t)^{-\lambda_1}$ for $t \in [T_1, T_2]$ and as $y^{(n-1)}$ is increasing (see (15)) we have $g(\Phi(t, y^{(n-1)}(t))) = g(y^{(n-1)}(t))$, $t \in [T_1, T_2]$. From this and from (9) and (18) we have

$$Z_m^{(n)}(t) \leq M_2 g(Z_m^{(n-1)}(t)), \quad t \in [T_1, T_2],$$

and hence,

$$\int_M^{2M} \frac{ds}{g(s)} \leq M_2(T_2 - T_1) \leq M_2(\tau - T).$$

The contradiction with (11) proves that (16) holds.

Further, we prove that

$$Z_m^{(n-1)}(t) < M_3(\tau - t)^{-\lambda_1}, \quad t \in [T, \tau_m], \quad (19)$$

holds for large m , say $m \geq \bar{m}_0$. If (19) is not valid, then either

(i) there exists $t_1 \in [T, \tau_m]$ such that

$$Z_m^{(n-1)}(t_1) = M_3(\tau - t_1)^{-\lambda_1}, \quad Z_m^{(n-1)}(\tau_m) \leq M_3(\tau - \tau_m)^{-\lambda_1}, \quad (20)$$

or

$$(ii) \quad Z_m^{(n-1)}(t) > M_3(\tau - t)^{-\lambda_1}, \quad (21)$$

in a left neighbourhood of $t = \tau_m$.

Let (i) be valid. It follows from (7), (13)–(16) that

$$Z_m^{(n)}(t) \geq M_1(Z_m^{(n-1)}(t))^\lambda, \quad t \in [t_1, \tau_m],$$

and the integration on $[t_1, \tau_m]$, (8) and (20) yield

$$\frac{\tau_m - t_1}{M_3^{\lambda-1}} \geq [Z_m^{(n-1)}(t_1)]^{1-\lambda} - [Z_m^{(n-1)}(\tau_m)]^{1-\lambda} \geq M_1(\lambda - 1)(\tau_m - t_1).$$

The contradiction with the definition of M_3 shows that case (i) is impossible.

Let (ii) hold and let $t_1 \in [T, \tau_m]$ be such that

$$Z_m^{(n-1)}(t) > M_3(\tau - t)^{-\lambda_1} \quad \text{on } [t_1, \tau_m].$$

Then the Taylor series theorem and (13) yield

$$\begin{aligned} k = Z_m(\tau_m) &\geq \int_{t_1}^{\tau_m} \frac{(\tau_m - s)^{n-2}}{(n-2)!} Z_m^{(n-1)}(s) ds \geq \frac{M_3}{(n-2)!} \int_{t_1}^{\tau_m} (\tau_m - s)^{n-2} (\tau - s)^{-\lambda} ds \\ &= \frac{-m M_3}{(n-1)!} \int_{t_1}^{\tau_m} (\tau - s)^{n-\lambda_1} \frac{d}{ds} \left(\left(1 - \frac{1}{m(\tau - s)} \right)^{n-1} \right) ds \\ &\geq \frac{M_3 m}{(n-1)!} A \left(1 - \frac{1}{m(\tau - t_1)} \right)^{n-1} \rightarrow \infty, \quad \text{for } m \rightarrow \infty, \end{aligned}$$

where

$$\begin{aligned} A &= (\tau - t_1)^{n-\lambda_1} \quad \text{for } n - \lambda_1 < 0, \\ A &= m^{\lambda_1-n} \quad \text{for } 0 \leq n - \lambda_1 < 1, \end{aligned}$$

as $n - 1 - \lambda_1 < 0$. Hence, (19) holds and the other derivatives of Z_m may be estimated.

It follows from the Taylor series theorem and from (11) that

$$\begin{aligned} M \leq Z_m^{(i)}(t) &= M \sum_{j=0}^{n-i-2} \frac{(t-T)^j}{j!} + \int_T^t \frac{(t-s)^{n-i-2}}{(n-i-2)!} Z_m^{(n-1)}(s) ds \\ &\leq M(n-1) + M_3 \int_T^t \frac{(\tau-s)^{n-i-2-\lambda_1}}{(n-i-2)!} \leq \varphi(t), \quad t \in J_m. \end{aligned} \quad (22)$$

From this, from (16) and (19), $\{Z_m^{(j)}\}$, $j = 0, 1, \dots, n-1$, $m = \bar{m}_0, \bar{m}_0 + 1, \dots$, are uniformly bounded with respect to j and m and hence, according to Arzela–Ascoli theorem (see [6, Lemma 10.2]), there exists a subsequence that converges uniformly to a solution y_k of (12). Moreover, the estimations (16), (19) and (22) hold also for y_k and hence $\Phi_0(t, y_k^{(i)}(t)) = y_k^{(i)}(t)$, $i = 0, 1, \dots, n-2$, $\Phi(t, y_k^{(n-1)}(t)) = y_k^{(n-1)}(t)$, $t \in J$. As (16), (19) and (22) do not depend on k and i , Arzela–Ascoli theorem yields the existence of a subsequence of $\{y_k\}_{k_0}^\infty$ that converges uniformly to a solution y of (1) fulfilling $\lim_{t \rightarrow \tau_-} y(t) = \infty$ and (4) holds.

Let $\bar{\lambda} = 1 + 1/(n-1)$. Let $\{\lambda_s\}_1^\infty$ be a sequence such that $\lambda_s > 1$, $\lim_{s \rightarrow \infty} \lambda_s = \bar{\lambda}$. Note, that according to (9) and (11) T is independent on the parametr λ and hence, there exist a solution y_s of (1), (4) with $\lambda = \lambda_s$, $s \in \{1, 2, \dots\}$ defined on $[T, \tau)$. Moreover, denote by $M_3(\lambda_s)$ the number M_3 from (9) for $\lambda = \lambda_s$. It is clear that

$$\bar{M}_3 = \sup_{s \in \{1, 2, \dots\}} M_3(\lambda_s) < \infty.$$

If M_3 is replaced by \bar{M}_3 , the estimations (16), (19) and (22) are valid for $Z_m = y_s$, $s = 1, 2, \dots$, too, and moreover, they do not depend on s . Hence, there exists Φ such that

$$|y_s^{(i)}(t)| \leq \Phi(t), \quad t \in [T, \tau), \quad i = 0, 1, \dots, n-1, \quad s = 1, 2, \dots,$$

and according to Arzela–Ascoli theorem, there exists a subsequence of $\{y_s\}_1^\infty$ that converges uniformly to a solution of (1), fulfilling (4) with $\lambda = \bar{\lambda}$. \square

Corollary 1. Let $\tau \in (0, \infty)$, $\lambda > 1$, $I = [\bar{\tau}, \tau] \subset R_+$, $M \in (0, \infty)$ and $M_1 \in (0, \infty)$ be such that

$$g(x) \geq |x|^\lambda \quad \text{for } |x| \geq M, \quad (23)$$

and (7) holds for $\alpha = 1$ and for $\alpha = -1$. Then there exists a singular solution y of (1) fulfilling (4) if and only if

$$1 < \lambda \leq 1 + \frac{1}{n-1}.$$

Proof. It follows from Theorem 1(ii) and Theorem 2. \square

Consider a special case of (1)

$$y^{(n)} = r(t)F(y)g(y^{(n-1)}), \quad (24)$$

where $\lambda > 0$, $r \in C^0(R_+)$, $F \in C^0(R)$, $g \in C^0(R)$ and let $M \in (0, \infty)$ and $\beta \in \{-1, 1\}$ be such that (23) and

$$\beta r(t) > 0 \quad \text{on } R_+, \quad F(x)x > 0 \quad \text{for } x \neq 0$$

hold.

Theorem A [2]. Let $c_0 \neq 0$. Then (23) has a singular solution y fulfilling (3) if

$$1 + \frac{1}{n-l-1} < \lambda \leq 1 + \frac{1}{n-l-2} \quad \text{for } l < n-2, \quad (25)$$

and

$$2 < \lambda \quad \text{for } l = n-2.$$

If, moreover, a constant $M_1 > 1$ exists such that $g(x) \leq M_1|x|^\lambda$ for $|x| \geq M$ then (23) has a singular solution y fulfilling (3) if and only if (25) is valid.

Applying Corollary 1 to (24) we obtain

Corollary 2. Let $\lambda > 1$, $\beta = 1$, $\tau \in (0, \infty)$ and $\liminf_{|x| \rightarrow \infty} |F(x)| > 0$. Then (24) has a singular solution y fulfilling (4) if and only if

$$1 < \lambda < 1 + \frac{1}{n-1}. \quad (26)$$

Note, that the intervals (25) (for different l) and (26) are disjunct. Moreover, they altogether cover the interval $(1, \infty)$ in case $\beta = 1$ (the interval $(1 + 1/(n-1), \infty)$ in case $\beta = -1$).

As a singular solution y of (24) is nonoscillatory if, and only if $\lim_{t \rightarrow \tau_-} |y^{(n-1)}(t)| = \infty$ (see, e.g., [1]), the following result follows from Corollary 2 and Theorem 1(i).

Corollary 3. Let $\liminf_{|x| \rightarrow \infty} |F(x)| > 0$.

- (i) If $\beta = 1$ and $\lambda > 1$, then (24) has a nonoscillatory singular solution.
- (ii) If $\beta = -1$, then (24) has a nonoscillatory singular solution if and only if $\lambda > 1 + 1/(n-1)$.

Note, that the result of Corollary 3(i) is known, see, e.g., [11, Theorem 11].

The existence of a singular solution fulfilling (4) in an open problem in case $\lambda \leq 1$ and $\beta = 1$. The following example shows that this situation is possible.

Example. Let $\tau \in (0, \infty)$. Then $y = 1/(\tau - t)$, $t \in [0, \tau)$ is a singular solution of $y''' = 3|y''| \operatorname{sgn} y$, that fulfills (4).

References

- [1] M. Bartušek, On existence of singular solutions of n th order differential equations, *Arch. Math. (Brno)* 36 (2000) 395–404.
- [2] M. Bartušek, J. Osička, On existence of singular solutions, *Georgian Math. J.* 8 (2001) 669–681.
- [3] T.A. Chanturia, On existence of singular and unbounded oscillatory solutions of differential equations of Emden–Fowler type, *Differentsial'nye Uravneniya* 28 (1992) 1009–1022 (in Russian).
- [4] C.V. Coffman, D.I. Ullrych, On the continuation of solutions of a certain non-linear differential equation, *Monatsh. Math. B* 71 (1967) 385–392.
- [5] C.V. Coffman, J.S.W. Wong, Oscillation and nonoscillation theorems for second order differential equations, *Funkcial. Ekvac.* 15 (1972) 119–130.
- [6] D.V. Izjumova, I.T. Kiguradze, Some remarks on solutions of $u'' + a(t)f(u) = 0$, *Differentsial'nye Uravneniya* 4 (1968) 589–605 (in Russian).
- [7] J. Jaroš, T. Kusano, On black hole solutions of second order differential equations with a singularity in the differential operator, *Funkcial. Ekvac.* 43 (2000) 491–509.
- [8] P. Hartman, *Ordinary Differential Equations*, Wiley, New York, 1964.
- [9] I.T. Kiguradze, Asymptotic properties of solutions of a nonlinear differential equation of Emden–Fowler type, *Izv. Akad. Nauk SSSR Ser. Mat.* 29 (1965) 965–986.
- [10] I.T. Kiguradze, Remark on oscillatory solutions of $u'' + a(t)|u|^n \operatorname{sign} u = 0$, *Čas. Pěst. Mat.* 92 (1967) 343–350.
- [11] I.T. Kiguradze, T. Chanturia, *Asymptotic Properties of Solution of Nonautonomous Ordinary Differential Equations*, Kluwer Academic, Dordrecht, 1993.